arXiv:1204.2921v1 [math.FA] 13 Apr 2012

OSTROWSKI TYPE INEQUALITIES VIA $h-$CONVEX FUNCTIONS WITH APPLICATIONS FOR SPECIAL MEANS AND P.D.F.'S

MEVLÜT TUNÇ

Abstract. In this paper, we establish some new Ostrowski type inequalities for the class of $h-$convex functions which are super-multiplicative or super-additive and nonnegative. Some applications for special means and PDF's are given.

1. **Introduction**

Let $f:I\subset[0,\infty)\rightarrow\mathbb{R}$ be a differentiable mapping on I° , the interior of the interval I, such that $f'\in L[a,b]$, where $a,b\in I$ with $a<b$. If $|f'(x)|\leq M$, then the following inequality:

$$\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right|\leq\frac{M}{b-a}\left[\frac{(x-a)^2+(b-x)^2}{2}\right] \tag{1.1}$$

holds. This result is known in the literature as the Ostrowski inequality. For recent results and generalizations concerning Ostrowski's inequality, see [10-15] and the references therein.

Definition 1. *[4] We say that $f:I\rightarrow\mathbb{R}$ is Godunova-Levin function or that f belongs to the class $Q(I)$ if f is non-negative and for all $x,y\in I$ and $t\in(0,1)$ we have*

$$f(tx+(1-t)y)\leq\frac{f(x)}{t}+\frac{f(y)}{1-t}. \tag{1.2}$$

Definition 2. *[1] We say that $f:I\subseteq\mathbb{R}\rightarrow\mathbb{R}$ is a $P-$function or that f belongs to the class $P(I)$ if f is nonnegative and for all $x,y\in I$ and $t\in[0,1]$, we have*

$$f(tx+(1-t)y)\leq f(x)+f(y). \tag{1.3}$$

Definition 3. *[3] Let $s\in(0,1]$. A function $f:(0,\infty]\rightarrow[0,\infty]$ is said to be $s-$convex in the second sense if*

$$f(tx+(1-t)y)\leq t^s f(x)+(1-t)^s f(y), \tag{1.4}$$

for all $x,y\in(0,b]$ and $t\in[0,1]$. This class of $s-$convex functions is usually denoted by K_s^2.

2000 *Mathematics Subject Classification.* 26D10, 26A15, 26A16, 26A51.

Key words and phrases. Ostrowski's inequality, $h-$convex, super-multiplicative, super-additive.

Definition 4. *[9] Let* $h : J \subseteq \mathbb{R} \to \mathbb{R}$ *be a positive function . We say that* $f : I \subseteq \mathbb{R} \to \mathbb{R}$ *is* $h-$*convex function, or that* f *belongs to the class* $SX(h, I)$*, if* f *is nonnegative and for all* $x, y \in I$ *and* $t \in [0, 1]$ *we have*

$$f(tx + (1 - t)y) \leq h(t) f(x) + h(1 - t) f(y). \tag{1.5}$$

If inequality (1.5) is reversed, then f is said to be $h-$concave, i.e. $f \in SV(h, I)$. Obviously, if $h(t) = t$, then all nonnegative convex functions belong to $SX(h, I)$ and all nonnegative concave functions belong to $SV(h, I)$; if $h(t) = \frac{1}{t}$, then $SX(h, I) = Q(I)$; if $h(t) = 1$, then $SX(h, I) \supseteq P(I)$; and if $h(t) = t^s$, where $s \in (0, 1)$, then $SX(h, I) \supseteq K_s^2$.

Remark 1. *[9]Let* h *be a non-negative function such that*

$$h(\alpha) \geq \alpha \tag{1.6}$$

for all $\alpha \in (0, 1)$*. For example, the function* $h_k(x) = x^k$ *where* $k \leq 1$ *and* $x > 0$ *has that property. If* f *is a non-negative convex function on* I *, then for* $x, y \in I$ *,* $\alpha \in (0, 1)$ *we have*

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y) \leq h(\alpha)f(x) + h(1 - \alpha)f(y). \tag{1.7}$$

So, $f \in SX(h, I)$*. Similarly, if the function* h *has the property:* $h(\alpha) \leq \alpha$ *for all* $\alpha \in (0, 1)$*, then any non-negative concave function* f *belongs to the class* $SV(h, I)$*.*

Definition 5. *[9]A function* $h : J \to \mathbb{R}$ *is said to be a super-multiplicative function if*

$$h(xy) \geq h(x) h(y) \tag{1.8}$$

for all $x, y \in J$*.*

If inequality (1.8) is reversed, then h is said to be a sub-multiplicative function. If equality is held in (1.8), then h is said to be a multiplicative function.

Definition 6. *[2]A function* $h : J \to \mathbb{R}$ *is said to be a super-additive function if*

$$h(x + y) \geq h(x) + h(y) \tag{1.9}$$

for all $x, y \in J$*.*

For recent results and generalizations concerning $h-$convex functions see [5-9] and references therein.

In [8] M.Z Sarıkaya, A. Sağlam and H. Yıldırım established the following Hadamard type inequality for $h-$convex functions:

Theorem 1. *[8] Let* $f\in SX(h, I)$*,* $a, b \in I$ *and* $f \in L_1([a, b])$*, then*

$$\frac{1}{2h\left(\frac{1}{2}\right)} f\left(\frac{a + b}{2}\right) \leq \frac{1}{b - a} \int_a^b f(x)\, dx \leq [f(a) + f(b)] \int_0^1 h(t)\, dt. \tag{1.10}$$

For recent results and generalizations concerning $h-$convex functions see [5-9].

The aim of this study is to establish some Ostrowski type inequalities for the class of functions whose derivatives in absolute value are $h-$convex and $h-$concave functions.

2. Ostrowski type inequalities for h−convex functions

In order to prove our main theorems, we need the following lemma that has been obtained in [13]:

Lemma 1. *[13]Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I^0 where $a, b \in I$ with $a < b$. If $f' \in L[a,b]$, then the following equality holds;*

$$\begin{aligned} & f(x) - \frac{1}{b-a}\int_a^b f(u)\,du \\ = {} & \frac{(x-a)^2}{b-a}\int_0^1 tf'(tx+(1-t)a)\,dt - \frac{(b-x)^2}{b-a}\int_0^1 tf'(tx+(1-t)b)\,dt \end{aligned}$$

for each $x \in [a,b]$.

Theorem 2. *Let $h : J \subseteq \mathbb{R} \to \mathbb{R}$ be a non-negative and super-multiplicative functions, $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I^0 such that $f' \in L[a,b]$, where $a, b \in I$ with $a < b$, and $h(\alpha) \geq \alpha$. If $|f'|$ is $h-$convex function on I and $|f'(x)| \leq M$, $x \in [a,b]$, then we have;*

$$\left| f(x) - \frac{1}{b-a}\int_a^b f(u)\,du \right| \leq \frac{M\left[(x-a)^2+(b-x)^2\right]}{b-a}\int_0^1 \left[h\left(t^2\right)+h\left(t-t^2\right)\right]dt. \tag{2.1}$$

for each $x \in [a,b]$.

Proof. By Lemma 1 and since $|f'|$ is $h-$convex, then we can write;

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a}\int_a^b f(u)\,du \right| \\ \leq {} & \frac{(x-a)^2}{b-a}\int_0^1 t\left|f'(tx+(1-t)a)\right|dt + \frac{(b-x)^2}{b-a}\int_0^1 t\left|f'(tx+(1-t)b)\right|dt \\ \leq {} & \frac{(x-a)^2}{b-a}\int_0^1 t\left[h(t)\left|f'(x)\right| + h(1-t)\left|f'(a)\right|\right]dt \\ & + \frac{(b-x)^2}{b-a}\int_0^1 t\left[h(t)\left|f'(x)\right| + h(1-t)\left|f'(b)\right|\right]dt \\ \leq {} & \frac{M(x-a)^2}{b-a}\int_0^1 \left[h^2(t)+h(t)h(1-t)\right]dt \\ & + \frac{M(b-x)^2}{b-a}\int_0^1 \left[h^2(t)+h(t)h(1-t)\right]dt \\ \leq {} & \frac{M\left[(x-a)^2+(b-x)^2\right]}{b-a}\int_0^1 \left[h\left(t^2\right)+h\left(t-t^2\right)\right]dt. \end{aligned}$$

The proof is completed. □

Remark 2. *If for* (2.1) *we choose $h(t) = t$, inequality* (2.1) *reduces to (1.1).*

In the next corollary, we will also make use of the Beta function of Euler type, which is for $x, y > 0$ defined as

$$\beta(x,y)=\int_0^1 t^{x-1}(1-t)^{y-1}\,dt=\frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

Corollary 1. *If for* (2.1) *we choose* $h(t)=t^s$*, then we have*

$$\begin{aligned}
\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right| &\leq \frac{M\left[(x-a)^2+(b-x)^2\right]}{b-a}\int_0^1\left[t^{2s}+\left(t-t^2\right)^s\right]dt\\
&= \frac{M\left[(x-a)^2+(b-x)^2\right]}{b-a}\int_0^1\left[t^{2s}+t^s(1-t)^s\right]dt\\
&= \frac{M\left[(x-a)^2+(b-x)^2\right]}{b-a}\left[\frac{1}{2s+1}+\frac{\Gamma(s+1)\Gamma(s+1)}{\Gamma(2s+2)}\right]\\
&= \frac{M\left[(x-a)^2+(b-x)^2\right]}{b-a}\left[\frac{\Gamma(2s+1)+s^2(\Gamma(s))^2}{(2s+1)\Gamma(2s+1)}\right]
\end{aligned}$$

The corresponding version for powers of the absolute value of the first derivative is incorporated in the following result:

Theorem 3. *Let* $h: J\subseteq\mathbb{R}\to\mathbb{R}$ *be a non-negative and superadditive functions,* $f: I\subseteq\mathbb{R}\to\mathbb{R}$ *be a differentiable mapping on* I^0 *such that* $f'\in L[a,b]$*, where* $a,b\in I$ *with* $a<b$*. If* $|f'|^q$ *is* h*-convex function on* $[a,b]$, $p,q>1$, $\frac{1}{p}+\frac{1}{q}=1$, $h(t)\geq t$ *and* $|f'(x)|\leq M$, $x\in[a,b]$*, then*

(2.2)

$$\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right|\leq\frac{Mh^{\frac{1}{q}}(1)}{b-a}\left(\int_0^1\left(h\left(t^p\right)dt\right)^{\frac{1}{p}}\right)\left((x-a)^2+(b-x)^2\right)$$

for each $x\in[a,b]$.

Proof. Suppose that $p>1$. From Lemma 1 and using the Hölder's inequality, we can write

$$\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right|\\
\leq\ &\frac{(x-a)^2}{b-a}\int_0^1 t\left|f'(tx+(1-t)a)\right|dt+\frac{(b-x)^2}{b-a}\int_0^1 t\left|f'(tx+(1-t)b)\right|dt\\
\leq\ &\frac{(x-a)^2}{b-a}\left(\int_0^1 t^p dt\right)^{\frac{1}{p}}.\left(\int_0^1\left|f'(tx+(1-t)a)\right|^q dt\right)^{\frac{1}{q}}\\
&+\frac{(b-x)^2}{b-a}\left(\int_0^1 t^p dt\right)^{\frac{1}{p}}.\left(\int_0^1\left|f'(tx+(1-t)b)\right|^q dt\right)^{\frac{1}{q}}.
\end{aligned}$$

Since $|f'|^q$ is $h-$convex and by using properties of $h-$convexity in the assumptions,

$$\begin{aligned}\int_0^1 \left|f'\left(tx+(1-t)a\right)\right|^q dt &\leq \int_0^1 \left[h(t)\left|f'(x)\right|^q + h(1-t)\left|f'(a)\right|^q\right] dt \\ &\leq M^q \int_0^1 \left[h(t)+h(1-t)\right] dt \\ &\leq M^q \int_0^1 h(1)\, dt = M^q h(1).\end{aligned}$$

Similarly, we can show that

$$\begin{aligned}\int_0^1 \left|f'\left(tx+(1-t)b\right)\right|^q dt &\leq \int_0^1 \left[h(t)\left|f'(x)\right|^q + h(1-t)\left|f'(b)\right|^q\right] dt \\ &\leq M^q h(1),\end{aligned}$$

and

$$\int_0^1 t^p dt \leq \int_0^1 h(t^p)\, dt.$$

Therefore, we obtain

$$\begin{aligned}\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right| &\leq Mh^{\frac{1}{q}}(1)\frac{(x-a)^2}{b-a}\left(\int_0^1 h(t^p)\,dt\right)^{\frac{1}{p}} \\ &\quad + Mh^{\frac{1}{q}}(1)\frac{(b-x)^2}{b-a}\left(\int_0^1 h(t^p)\,dt\right)^{\frac{1}{p}} \\ &= \frac{Mh^{\frac{1}{q}}(1)}{b-a}\left(\int_0^1 h(t^p)\,dt\right)^{\frac{1}{p}}\left((x-a)^2+(b-x)^2\right)\end{aligned}$$

The proof is completed. □

Corollary 2. *If for* (2.2) *we choose* $h(t)=t$*, then we have*

$$\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \leq \frac{M}{b-a}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left((x-a)^2+(b-x)^2\right). \tag{2.3}$$

Remark 3. *Since* $\left(\frac{1}{p+1}\right)^{\frac{1}{p}} > \frac{1}{2}$ *for any* $p>1$*, then we observe that the inequality (2.3) is better than the inequality (1.1) meaning that the approach via the Hölder's inequality is a better approach than that through classical inequality.*

A different approach leads to the following result.

Theorem 4. *Let* $h: J\subseteq\mathbb{R}\to\mathbb{R}$ *be a non-negative and supermultiplicative functions,* $f: I\subseteq\mathbb{R}\to\mathbb{R}$ *be a differentiable mapping on* I^0 *such that* $f'\in L[a,b]$*, where* $a,b\in I$ *with* $a<b$*. If* $|f'|^q$ *is* $h-$*convex function on* $[a,b]$, $q>1$, $h(\alpha)\geq\alpha$ *and* $|f'(x)|\leq M$, $x\in[a,b]$*, then*

$$\begin{aligned}&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\ &\leq \frac{\sqrt[q]{2}M}{2(b-a)}\left((x-a)^2+(b-x)^2\right)\left(\int_0^1 \left(h\left(t^2\right)+h\left(t-t^2\right)dt\right)^{\frac{1}{q}}\right)\end{aligned} \tag{2.4}$$

for each $x \in [a,b]$.

Proof. Suppose that $q \geq 1$. From Lemma 1 and using the power mean inequality , we have

$$\begin{aligned}
&\left| f(x) - \frac{1}{b-a}\int_a^b f(u)\,du \right| \\
\leq\ & \frac{(x-a)^2}{b-a}\int_0^1 t\left|f'(tx+(1-t)a)\right|dt + \frac{(b-x)^2}{b-a}\int_0^1 t\left|f'(tx+(1-t)b)\right|dt \\
\leq\ & \frac{(x-a)^2}{b-a}\left(\int_0^1 t^p dt\right)^{1-\frac{1}{q}} . \left(\int_0^1 t\left|f'(tx+(1-t)a)\right|^q dt\right)^{\frac{1}{q}} \\
& + \frac{(b-x)^2}{b-a}\left(\int_0^1 t^p dt\right)^{1-\frac{1}{q}} . \left(\int_0^1 t\left|f'(tx+(1-t)b)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}$$

Since $|f'|^q$ is $h-$convex, we have

$$\begin{aligned}
\int_0^1 t\left|f'(tx+(1-t)a)\right|^q dt &\leq \int_0^1 \left[th(t)\left|f'(x)\right|^q + th(1-t)\left|f'(a)\right|^q\right]dt \\
&\leq \left|f'(x)\right|^q \int_0^1 h(t)h(t)\,dt + \left|f'(a)\right|^q \int_0^1 h(t)h(1-t)\,dt \\
&\leq M^q\left[\int_0^1 h\left(t^2\right)dt + \int_0^1 h\left(t-t^2\right)dt\right].
\end{aligned}$$

Similarly, we can observe that

$$\begin{aligned}
\int_0^1 t\left|f'(tx+(1-t)b)\right|^q dt &\leq \left|f'(x)\right|^q \int_0^1 h(t)h(t)\,dt + \left|f'(b)\right|^q \int_0^1 h(t)h(1-t)\,dt \\
&\leq M^q\left\{\int_0^1 h\left(t^2\right)dt + \int_0^1 h\left(t-t^2\right)dt\right\}.
\end{aligned}$$

Therefore, we deduce

$$\begin{aligned}
&\left| f(x) - \frac{1}{b-a}\int_a^b f(u)\,du \right| \\
\leq\ & \frac{(x-a)^2}{b-a}\left(\frac{1}{2}\right)^{1-\frac{1}{q}}\left(M^q\int_0^1\left(h\left(t^2\right)+h\left(t-t^2\right)\right)dt\right)^{\frac{1}{q}} \\
& + \frac{(b-x)^2}{b-a}\left(\frac{1}{2}\right)^{1-\frac{1}{q}}\left(M^q\int_0^1\left(h\left(t^2\right)+h\left(t-t^2\right)\right)dt\right)^{\frac{1}{q}} \\
=\ & M\left(\frac{1}{2}\right)^{1-\frac{1}{q}}\left(\int_0^1\left(h\left(t^2\right)+h\left(t-t^2\right)\right)dt\right)^{\frac{1}{q}}\left(\frac{(x-a)^2+(b-x)^2}{(b-a)}\right) \\
=\ & \sqrt[q]{2}M\left(\int_0^1\left(h\left(t^2\right)+h\left(t-t^2\right)\right)dt\right)^{\frac{1}{q}}\left(\frac{(x-a)^2+(b-x)^2}{2(b-a)}\right)
\end{aligned}$$

and the proof is completed. □

Remark 4. *i) In the previous inequalities, one can obtain several midpoint type inequalities by setting* $x = \frac{a+b}{2}$.

ii) All of the above inequalities obviously hold for non-negative convex functions. Simply choose $h(t)=t$, $h(t)=\frac{1}{t}$, $h(t)=1$ *and* $h(t)=t^s$ *in each of those results, to get desired results respectively for non-negative convex functions,* $Q(I)$, $P(I)$ *and* K_s^2.

The following result holds for $h-$concave functions.

Theorem 5. *Let* $h: J\subseteq\mathbb{R}\to\mathbb{R}$ *be a non-negative and superadditive functions,* $f: I\subseteq\mathbb{R}\to\mathbb{R}$ *be a differentiable mapping on* I^0 *such that* $f'\in L[a,b]$ *,where* $a,b\in I$ *with* $a<b$. *If* $|f'|^q$ *is* $h-$*concave function on* $[a,b]$, $p,q>1$, $\frac{1}{p}+\frac{1}{q}=1$, $h(t)\geq t$, *then*

$$\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \leq \frac{1}{\sqrt[q]{2}\,(p+1)^{\frac{1}{p}}\,h^{\frac{1}{q}}\left(\frac{1}{2}\right)}\left[\frac{(x-a)^2}{b-a}\left|f'\left(\frac{x+a}{2}\right)\right|+\frac{(b-x)^2}{b-a}\left|f'\left(\frac{x+b}{2}\right)\right|\right] \tag{2.5}$$

for each $x\in[a,b]$.

Proof. Suppose that $p>1$. From Lemma 1 and using the Hölder's inequality, we can write

$$\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{(x-a)^2}{b-a}\int_0^1 t\,|f'(tx+(1-t)a)|\,dt+\frac{(b-x)^2}{b-a}\int_0^1 t\,|f'(tx+(1-t)b)|\,dt \\
\leq\ & \frac{(x-a)^2}{b-a}\left(\int_0^1 t^p dt\right)^{\frac{1}{p}}.\left(\int_0^1 |f'(tx+(1-t)a)|^q\,dt\right)^{\frac{1}{q}} \\
&+\frac{(b-x)^2}{b-a}\left(\int_0^1 t^p dt\right)^{\frac{1}{p}}.\left(\int_0^1 |f'(tx+(1-t)b)|^q\,dt\right)^{\frac{1}{q}}.
\end{aligned} \tag{2.6}$$

But since $|f'|^q$ is $h-$concave, using the inequality (1.10), we have

$$\left(\int_0^1 |f'(tx+(1-t)a)|^q\,dt\right)\leq\frac{1}{2h\left(\frac{1}{2}\right)}\left|f'\left(\frac{x+a}{2}\right)\right|^q \tag{2.7}$$

and

$$\left(\int_0^1 |f'(tx+(1-t)b)|^q\,dt\right)\leq\frac{1}{2h\left(\frac{1}{2}\right)}\left|f'\left(\frac{x+b}{2}\right)\right|^q. \tag{2.8}$$

By combining the above numbered inequalities, we obtain

$$
\begin{aligned}
&\left|f(x)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{(x-a)^2}{b-a}\frac{1}{(p+1)^{\frac{1}{p}}}\left(\frac{1}{2h\left(\frac{1}{2}\right)}\right)^{\frac{1}{q}}\left|f'\left(\frac{x+a}{2}\right)\right| \\
&+\frac{(b-x)^2}{b-a}\frac{1}{(p+1)^{\frac{1}{p}}}\left(\frac{1}{2h\left(\frac{1}{2}\right)}\right)^{\frac{1}{q}}\left|f'\left(\frac{x+b}{2}\right)\right| \\
=\ & \frac{1}{\sqrt[q]{2}\,(p+1)^{\frac{1}{p}}\,h^{\frac{1}{q}}\left(\frac{1}{2}\right)}\left[\frac{(x-a)^2}{b-a}\left|f'\left(\frac{x+a}{2}\right)\right|+\frac{(b-x)^2}{b-a}\left|f'\left(\frac{x+b}{2}\right)\right|\right]
\end{aligned}
$$

The proof is completed. □

A midpoint type inequality for fuctions whose derivetives in absolute value are $h-$concave may be obtained from the previous result as follows:

Corollary 3. *If for* (2.5) *we choose* $x=\frac{a+b}{2}$, *then we get*

$$
\begin{aligned}
&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{b-a}{\sqrt[q]{2^{2q+1}}\,(p+1)^{\frac{1}{p}}\,h^{\frac{1}{q}}\left(\frac{1}{2}\right)}\left[\left|f'\left(\frac{3a+b}{4}\right)\right|+\left|f'\left(\frac{a+3b}{4}\right)\right|\right].
\end{aligned}
\tag{2.9}
$$

For instance, if $h(t)=t$, *then we obtain*

$$
\begin{aligned}
&\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(u)\,du\right| \\
\leq\ & \frac{b-a}{\sqrt[q]{2^{2q+2}}\,(p+1)^{\frac{1}{p}}}\left[\left|f'\left(\frac{3a+b}{4}\right)\right|+\left|f'\left(\frac{a+3b}{4}\right)\right|\right].
\end{aligned}
\tag{2.10}
$$

where $|f'|^q$ *is* $h-$*concave function on* $[a,b]$, $p,q>1$.

3. Applications to special means

We consider the means for arbitrary positive numbers a,b $(a\neq b)$ as follows;
The arithmetic mean

$$A(a,b)=\frac{a+b}{2}$$

The generalized log-mean :

$$L_p(a,b)=\left[\frac{b^{p+1}-a^{p+1}}{(p+1)(b-a)}\right]^{\frac{1}{p}},\ p\in\mathbb{R}\setminus\{-1,0\}.$$

The identric mean:

$$I(a,b)=\frac{1}{e}\left(\frac{b^b}{a^a}\right)^{\frac{1}{b-a}}$$

Now, using the result of Section 2, we give some applications to special means of real numbers.

In [9], the following example is given:

Let h be a function defined by $h(x) = (c+x)^{p-1}$, $x \geq 0$. If $c = 0$, then the function h is multiplicative. If $c \geq 1$, then for $p \in (0,1)$ the function h is super-multiplicative and for $p > 1$ the function h is sub-multiplicative.

Hence, for $c = 1$, $p \in (0,1)$, we have $h(t) = (1+t)^{p-1}$, $t \geq 0$ is supermultiplicative. Let $f(x) = x^n$, $x > 0$, $|n| \geq 2$ is $h-$convex functions.

Proposition 1. *Let* $0 < a < b$, $p \in (0,1)$ *and* $|n| \geq 2$. *Then*

$$|A^n(a,b) - L_n^n(a,b)| \leq \frac{M(b-a)}{4}\left[\int_0^1 \left(1+t^2\right)^{p-1} dt + \int_0^1 \left(1+t-t^2\right)^{p-1} dt\right]$$

Proof. The inequality is derived from (2.1) with $x = \frac{a+b}{2}$ applied to the $h-$convex functions $f : \mathbb{R} \to \mathbb{R}$, $f(x) = x^n$, $|n| \geq 2$ and $h : \mathbb{R} \to \mathbb{R}$, $h(t) = (1+t)^{p-1}$, $p \in (0,1)$. The details are disregarded. ☐

Proposition 2. *Let* $0 < a < b$, $p \in (0,1)$, $q > 1$ *and* $|n| \geq 2$. *Then*

$$|A^n(a,b) - L_n^n(a,b)| \leq \frac{\sqrt[p]{2}M(b-a)}{8}\left[\int_0^1 \left(1+t^2\right)^{p-1} dt + \int_0^1 \left(1+t-t^2\right)^{p-1} dt\right]^{\frac{1}{q}}$$

Proof. The inequality is derived from (2.3) with $x = \frac{a+b}{2}$ applied to the $h-$convex functions $f : \mathbb{R} \to \mathbb{R}$, $f(x) = x^n$, $|n| \geq 2$ and $h : \mathbb{R} \to \mathbb{R}$, $h(t) = (1+t)^{p-1}$, $p \in (0,1)$. The details are disregarded. ☐

Proposition 3. *Let* $0 < a < b$ *and* $p, q > 1$. *Then we have*

$$\begin{aligned} &|\ln(A(a,b)+1) - (b-a)\ln I(a+1,b+1)| \\ \leq\ & \frac{b-a}{\sqrt[q]{2^{2q+2}}\,(p+1)^{\frac{1}{p}}}\left[\frac{1}{3a+b+4} + \frac{1}{a+3b+4}\right]. \end{aligned} \tag{3.1}$$

Proof. The inequality is derived from (2.10) applied to the $h-$concave function $f : [a,b] \to \mathbb{R}$, $f(x) = \ln(x+1)$. The details are disregarded. ☐

4. **Applications for P.D.F.'s**

Let X be a random variable taking values in the finite interval $[a,b]$, with the probability density function $f : [a,b] \to [0,1]$ with the cumulative distribution function $F(x) = \Pr(X \leq x) = \int_a^b f(t)dt$.

Theorem 6. *Under the assumptions of Theorem 2, we have the inequality;*

$$\begin{aligned} &\left|f(x) - \frac{1}{b-a}(b - E(x))\right| \\ &\leq \frac{M\left[(x-a)^2 + (b-x)^2\right]}{(b-a)} \int_0^1 \left[h\left(t^2\right) + h(t-t^2)\right] dt \end{aligned}$$

where $E(x)$ *is the expectation of* X.

Proof. The proof is immediate follows from the fact that;

$$E(x)=\int_a^b t dF(t)=b-\int_a^b F\left(t\right)dt.$$

□

Theorem 7. *Under the assumptions of Theorem 3, we have the inequality;*

$$\begin{aligned}&\left|f\left(x\right)-\frac{1}{b-a}\left(b-E(x)\right)\right|\\&\leq\frac{Mh^{\frac{1}{q}}\left(1\right)}{b-a}\left(\left(x-a\right)^2+\left(b-x\right)^2\right)\int_0^1\left(h\left(t^p\right)\right)^{\frac{1}{p}}dt\end{aligned}$$

where $E(x)$ is the expectation of X.

Proof. Likewise the proof of the previous theorem, by using the fact that;

$$E(x)=\int_a^b t dF(t)=b-\int_a^b F\left(t\right)dt$$

the proof is completed. □

Theorem 8. *Under the assumptions of Theorem 4, we have inequality;*

$$\begin{aligned}&\left|f\left(x\right)-\frac{1}{b-a}\left(b-E(x)\right)\right|\\&\leq\frac{\sqrt[q]{2}M}{2\left(b-a\right)}\left(\left(x-a\right)^2+\left(b-x\right)^2\right)\int_0^1\left(\left[h\left(t^2\right)+h(t-t^2)\right]\right)^{\frac{1}{q}}dt\end{aligned}$$

where $E(x)$ is the expectation of X.

Proof. The proof is similar to the previous theorem. □

Acknowledgement 1. *The author gives his warm thanks to the Editor and the Authors for their precious papers in the reference list.*

References

[1] Dragomir, S.S., Pečarić, J. and Persson, L.E. *Some inequalities of Hadamard type*, Soochow J.Math., 21, 335–241, 1995.

[2] Alzer, H. *A superadditive property of Hadamard's gamma function*, Abh. Math. Semin. Univ. Hambg., 79 (2009), 11-23.

[3] Hudzik, H., Maligranda, L. *Some remarks on $s-$convex functions*, Aequationes Math., 48, 100–111, 1994.

[4] Godunova, E.K., Levin, V.I. *Neravenstva dlja funkcii sirokogo klassa, soderzascego vypuklye, monotonnye i nekotorye drugie vidy funkii*, Vycislitel. Mat. i. Fiz. Mezvuzov. Sb. Nauc. Trudov, MGPI, Moskva, pp. 138–142, 1985.

[5] Bombardelli, M., Varošanec, S. *Properties of $h-$convex functions related to the Hermite–Hadamard–Fejér inequalities*, Comput. Math. Appl. 58 (2009) 1869–1877.

[6] Burai, P., Hazy, A. *On approximately $h-$convex functions,* J. Convex Anal. 18 (2) (2011) 447–454.

[7] Özdemir, M.E., Gürbüz, M. and Akdemir, A.O. *Inequalities for $h-$Convex Functions via Further Properties*, RGMIA Research Report Collection Volume 14, article 22, 2011.

[8] Sarıkaya, M.Z., Sağlam, A., Yıldırım, H. *On some Hadamard-type inequalities for $h-$convex functions,* J. Math. Inequal. 2 (3) (2008) 335–341.

[9] Varošanec, S. *On $h-$convexity*, J. Math. Anal. Appl., Volume 326, Issue 1, 303–311, 2007.

[10] Alomari, M., Darus, M. *Some Ostrowski type inequalities for convex functions with applications*, RGMIA 13 (1) (2010) article No. 3. Preprint.

[11] Alomari, M., Darus, M. *Some Ostrowski type inequalities for quasi-convex functions with applications to special means*, RGMIA 13 (2) (2010) article No. 3. Preprint.

[12] Alomari, M., Darus, M., Dragomir, S.S., Cerone, P. *Ostrowski type inequalities for functions whose derivatives are $s-$convex in the second sense*, Appl. Math. Lett. Volume 23 (2010) 1071-1076.

[13] Cerone, P., Dragomir, S.S. *Ostrowski type inequalities for functions whose derivatives satisfy certain convexity assumptions*, Demonstratio Math. 37 (2) (2004) 299 308.

[14] Barnett, N.S., Cerone, P., Dragomir, S.S., Pinheiro, M.R., Sofo, A. *Ostrowski type inequalities for functions whose modulus of derivatives are convex and applications*, RGMIA Res. Rep. Coll. 5 (2) (2002) Article 1. Online: http://rgmia.vu.edu.au/v5n2.html.

[15] Dragomir, S.S., Sofo, A. *Ostrowski type inequalities for functions whose derivatives are convex*, in: Proceedings of the 4th International Conference on Modelling and Simulation, November 11 13, 2002. Victoria University, Melbourne, Australia, RGMIA Res. Rep. Coll. 5 (2002) Supplement, Article 30. Online: http://rgmia.vu.edu.au/v5(E).html.

KILIS 7 ARALIK UNIVERSITY, FACULTY OF SCIENCE AND ARTS, DEPARTMENT OF MATHEMATICS, KILIS, 79000, TURKEY.

E-mail address: `mevluttunc@kilis.edu.tr`